arXiv:0806.3552v2 [nucl-th] 1 Jul 2008

Binding Energies and Radii of the Nuclei with $N \geq Z$ in an Alpha-Cluster Model

G. K. NIE

Institute of Nuclear Physics, Ulugbek, Tashkent 702132, Uzbekistan

---

Abstract

Using the surface tension energy put in dependence on the number of α-clusters in the core in a phenomenological model representing a nucleus as a core and a nuclear molecule on its surface leads to widening the number of isotopes to be described from the narrow strip of β-stability to the isotopes with $N \geq Z$. The number of alpha-clusters in the molecule is obtained from the analysis of experimental binding energies and the specific density of the core binding energy ρ and the radii are calculated. It is shown that for the isotopes of one Z with growing A the number of alpha-clusters of the molecule decreases mostly to 3 and ρ increases to reach a saturation value within $\rho = 2.5 \div 2.7$ MeV/fm^3 at the β - stable isotopes, so the narrow strip of the binding energies of the β-stable isotopes with $Z \leq 84$ is outlined by a function of one variable Z.

Key words: nuclear structure; alpha-cluster model; core; Coulomb energy; surface tension energy, binding energy; charge radius.
PACS: 21.60.-n; 21.60.Gx; 21.10.Dr; 21.60.Cs.

---

1 The formulas to calculate binding energies and radii

In the representation of a nucleus as a core and a nuclear molecule on the surface of the core the nuclear binding energies and the radii of β- stable isotopes are calculated [1]. The calculations are made in the framework of an α-cluster model based on pn-pair interactions with using isospin invariance of nuclear force [2,3]. In the framework of the model some parameters like the binding energy and the energy of Coulomb repulsion between two nearby clusters had been found to be $\epsilon_{\alpha\alpha} = 2.425$ MeV and $\epsilon^C_{\alpha\alpha} = 1.925$ MeV, so that the nuclear force energy is to be $\epsilon^{nuc}_{\alpha\alpha} = \epsilon_{\alpha\alpha} + \epsilon^C_{\alpha\alpha} = 4.350$ MeV [3] as well as

---

Email address: `galani@Uzsci.net` (G. K. NIE).

Preprint submitted to Elsevier *1 July 2008*

the energy of nuclear force of one α-cluster $\epsilon_\alpha^{nuc} = \epsilon_\alpha + \epsilon_\alpha^C = 29.060$ MeV [3,4] where $\epsilon_\alpha = E_{^4\mathrm{He}} = 28.296$ MeV and $\epsilon_\alpha^C = 0.764$ MeV.

One of the main findings [3] of the phenomenological model is the formula for binding energy E of the symmetrical nuclei $(N = Z, Z < 30)$ with the number of α - clusters $N_\alpha = Z/2$

$$E = N_\alpha \epsilon_\alpha + 3(N_\alpha - 2)\epsilon_{\alpha\alpha}, \tag{1}$$

for the odd $Z_1 = Z + 1$ nuclei the energy $E_1 = E + 13.9$ MeV. It is suggested that the number of short range links in a nucleus consisting of N_α α-clusters is $3(N_\alpha - 2)$ and that the long range part of the Coulomb interactions must be compensated with the surface tension energy E^{st}. Then with using isospin invariance of nuclear force the empirical values of the Coulomb energy, the energy of surface tension and the empirical values of the distance of the position of the last alpha-cluster R_α in the system of the center of the masses of the remote $N_\alpha - 4$ α-clusters have been obtained [3]. From the analysis of these values the formulas for Coulomb radius $R_C = 1.869 N_\alpha^{1/3}$fm (see (22) [3]), for the radius of the position of the last α-cluster $R_\alpha = 2.168(N_\alpha - 4)^{1/3}$ fm (see (21) [2,3]) have been found. The Coulomb energy of the charge sphere of the radius R_C is $E^C = 3/5 Z^2 e^2 / R_C$, which after simplifying is $1.848(N_\alpha)^{5/3}$ MeV [3]. The binding energy of the excess nn-pairs for the beta-stable nuclei $E_{\Delta N} = \sum_1^{N_{nn}} E_{i_{nn}}$ (see (13) [3]) where $E_{i_{nn}}$ is the binding energy of the i^{th} nn-pair in the core. The nn-pairs fill out the free space in the core which appears due to the difference between the charge and the matter radii of an alpha-cluster. Thus, the binding energy of all β - stable nuclei with an accuracy in a few MeV is calculated as the sum $E = E^{nuc} + E^{st} - E^C + E_{\Delta N}$ where E^{nuc} is the energy of nuclear force in short range links (see (7) [3]).

There is a clear relation between the nuclei $A(Z, N)$ and $A_1(Z_1, N + 2)$ where $Z_1 = Z + 1$, N is an even number and $N \geq Z$ [1,2,3]. Then $A_1 = A + 3$. The nuclei have equal cores with N_α^{core} α - clusters and with the same number of excess nn-pairs $N_{nn} = \Delta N/2$, where $\Delta N = N - Z$. The peripheral molecule on the surface of the core consists of N_α^{ml} α - clusters for the even nucleus and of $N_\alpha^{ml} + 0.5$ for the odd nucleus. In case of the nucleus $A_1(Z_1, N + 2)$ one neutron is stuck to the single pn-pair. It has been taken into account [1] that the number of short range links in the core has to be $3(N_\alpha^{core} - 2) + 6$, because the total number of links is $3(N_\alpha - 2) = 3(N_\alpha^{ml} - 2) + 3(N_\alpha^{core} - 2) + 6$. Therefore the energy of six short range links $6\epsilon_{\alpha\alpha} = 6(\epsilon_{\alpha\alpha}^{nuc} - \epsilon_{\alpha\alpha}^C)$ is added to the binding energy of N_α^{core} α - clusters in calculation of the core binding energy.

So the binding energies E and E_1 of the nuclei $A(Z, N)$ and $A_1(Z_1, N + 2)$ are calculated as follows, see (4,5)[1]

$$E = E_{N_\alpha^{ml}} - E^C_{N_\alpha^{ml} N_\alpha^{core}} + E_{core}; E_1 = E_{N_\alpha^{ml+0.5}} - E^C_{N_\alpha^{ml} N_\alpha^{core}} + E_{core}, \tag{2}$$

where $E_{N_\alpha^{ml}}$ and $E_{N_\alpha^{ml+0.5}}$ are the experimental binding energies of the nuclei with the total number of alpha-clusters equal to N_α^{ml} and $N_\alpha^{ml+0.5}$ (for example $E_3 = E_{^{12}\mathrm{C}}$ and $E_{3.5} = E_{^{15}\mathrm{N}}$), $E^C_{N_\alpha^{ml} N_\alpha^{core}}$ is the energy of the Coulomb interaction between the peripheral molecule and the core

$$E^C_{N_\alpha^{ml} N_\alpha^{core}} = 2N_\alpha^{ml} 2N_\alpha^{core} e^2/R_\alpha, \tag{3}$$

E_{core} is the core binding energy

$$E_{core} = E_{\Delta N} + E_{N_\alpha^{core}} + 6\epsilon_{\alpha\alpha}, \tag{4}$$

where the binding energy of excess nn-pairs $E_{\Delta N} = \sum_1^{N_{nn}} E_{i_{nn}}$ was approximated (see (12) in [1]) with the following formula in dependence on the number of excess nn-pairs N_{nn}

$$E_{\Delta N} = (21.93 - 0.762 N_{nn}^{2/3}) N_{nn}, \tag{5}$$

$E_{N_\alpha^{core}}$ stands for the binding energy of N_α^{core} of core α-clusters

$$E_{N_\alpha^{core}} = E^{nuc}_{N_\alpha^{core}} - E^C_{N_\alpha^{core}} + E^{st}, \tag{6}$$

where $E^{nuc}_{N_\alpha^{core}} = N_\alpha^{core} \epsilon_\alpha^{nuc} + 3(N_\alpha^{core} - 2)\epsilon^{nuc}_{\alpha\alpha}$, $E^C_{N_\alpha^{core}} = 1.848 (N_\alpha^{core})^{5/3}$; E^{st} is the surface tension energy

$$E^{st} = (N_\alpha + 1.7)(N_\alpha^{core})^{2/3}. \tag{7}$$

The original formula $E^{st} = (N_\alpha + 1.7)(N_\alpha - 4)^{2/3}$ (see (17) in Ref. [1]) was proposed in Ref. [3] as an approximation function to the factorized sum of the square radii of $N_\alpha - 4$ clusters for the nuclei with $Z \geq 30$. The part $(N_\alpha - 4)^{2/3}$ is changed here for $(N_\alpha^{core})^{2/3}$. In case when N_α^{ml}=4, eq. (7) and the original one coincide. So all the formulas are from [1] with one little change in the formula for the surface tension energy.

For the isotopes of the nuclei with $Z, Z_1 \leq 29$ the binding energy of the α-clusters is known from the energy of the symmetrical nuclei with $Z = N$ (1). So the binding energy for the isotopes is to be trivial $E = N\epsilon_\alpha + 3(N_\alpha - 2)\epsilon_{\alpha\alpha} + E_{\Delta N}$. The energy can be written in the terms of core and a peripheral molecule (2) with $E_{core} = N_\alpha^{core}\epsilon_\alpha + 3(N_\alpha^{core} - 2)\epsilon_{\alpha\alpha} + 6\epsilon_{\alpha\alpha} + E_{\Delta N} + E^C_{N_\alpha^{ml} N_\alpha^{core}}$.

The energy E (2) for the cases almost does not depend on N_α^{ml} varied within $2 \div 5$.

Two molecules on the surface of the core, let them consist of N_α^{ml1} and N_α^{ml2} alpha-clusters, are needed in describing the isotopes with a considerable deficiency of excess neutrons in the nuclei with big Z, see section 2. Then the binding energy is as follows [1]

$$E = E_{N_\alpha^{ml1}} + E_{N_\alpha^{ml2}} - (E^C_{N_\alpha^{ml1}(N_\alpha^{core}+N_\alpha^{ml2})} + E^C_{N_\alpha^{ml2}N_\alpha^{core}}) + E_{core}. \quad (8)$$

For the odd nucleus $A_1(Z_1, N+2)$ to calculate E_1 one pn-pair is added to one of the molecules. Here $E_{N_\alpha^{ml1}}$ is exchanged for $E_{N_\alpha^{ml1+0.5}}$. The value E_{core} is calculated by (4) with $N_\alpha^{core} = N_\alpha - (N_\alpha^{ml1} + N_\alpha^{ml2})$. Unlike [1] the total number of short links in the core does not depend on whether there is one or two molecules on the surface of the core.

The following formulas (9-12) have been proposed to estimate radii of the nuclei $A(Z, N)$ on $N_\alpha = Z/2$ and the odd nuclei $A_1(Z_1, N+2)$ on $N_\alpha + 0.5$ [1,2,3]. The simplest one for $Z, Z_1 \geq 24$ is the following

$$R = r_\alpha N_\alpha^{1/3}. \quad (9)$$

The core prevails in the cases, so the value r_α is to be the charge radius of a core α-cluster. To define r_α the empirical radii in [5], Table IIIA, with 116 data for the isotopes $A(Z, N)$ and $A_1(Z_1, N+2)$ with $Z \geq 24$ have been fitted. The authors analyze the muonic atom transition energies with using the same model for all observed there isotopes for the nuclei with $Z, Z_1 \leq 60$ and $77 < Z, Z_1 \leq 83$. The value $r_\alpha = 1.595$ fm has been obtained here with the rms deviation from the experimental data $\delta = 0.030$ fm. For the 147 isotopes $A(Z, N)$ and $A_1(Z_1, N+2)$ with $Z \geq 6$ of the table $\delta = 0.064$ fm.

The charge radius for the nuclei with $Z, Z_1 \geq 6$ can be estimated from adding the volumes of the charges of the core and the peripheral molecule [1]

$$R^3 = r^3_{^4He} N_\alpha^{ml} + r^3_\alpha N_\alpha^{core}, \quad (10)$$

where the radius of a peripheral alpha-cluster $r_{^4He}$=1.71 fm. For the nuclei without core, i.e. with $N_\alpha \leq 5$, N_α^{core}=0. Fitting the data of the Table IIIA [5] for the isotopes with $Z \geq 24$ with the values N_α^{ml}, obtained from analysis of binding energies, see section 2, ($N_\alpha^{ml} = 3$ in most of the cases), gives the value $r_\alpha = 1.574$ fm with $\delta = 0.033$ fm. As for the isotopes with $12 \leq Z, Z_1 \leq 23$ we tried N_α^{ml}= 5, which means the nuclear molecule ^{20}Ne for the even isotopes and ^{23}Na for the odd isotopes. Then for the 147 isotopes with $Z \geq 6$ of the table $\delta = 0.040$ fm.

Another way to calculate charge radii for the nuclei with $Z, Z_1 \geq 6$ is the following [2]

$$N_\alpha R^2 = N_\alpha^{core}(r_\alpha(N_\alpha^{core})^{1/3})^2 + N_\alpha^{ml} R_\alpha^2, \tag{11}$$

where the square core radius and the square distance of the position of the last α-cluster in the system of the center of mass of the core (for the $Z, Z_1 \leq 23$ the empirical values R_α are used, see (17) in [3]) are added with their weights to be equal to the square radius of the nucleus weighed with N_α. In the cases when $N_\alpha^{core} < N_\alpha^{ml}$ ($N_\alpha < 10$ and $N_\alpha^{ml} = 5$) the formula is rewritten for the center of mass of the peripheral molecule. Then N_α^{ml} and N_α^{core} exchange their places and r_α is replaced with $r_{^4He}$. The value $r_\alpha = 1.595$ fm obtained from fitting 116 data for the isotopes with $Z \geq 24$ of Table IIIA [5] gives $\delta = 0.038$ fm. For the 147 isotopes with $Z \geq 6$ of the table $\delta = 0.040$ fm.

The next formula estimates the radii for $Z, Z_1 \geq 6$ by adding the volumes of the charge of the peripheral N_α^{ml} α-clusters and the body of the core, consisting of the volume occupied by the bodies of the alpha-clusters and the volume of the excess nn-pairs [2]

$$R_m^3 = N_\alpha^{ml} r_{^4He}^3 + N_\alpha^{core}(4r_{p/n}^3) + N_{nn}(2r_n^3), \tag{12}$$

where $r_{p/n}$=0.954 fm stands for the radius of the volume which is one nucleon's share in the volume of an α-cluster body, r_n =0.796 fm is the radius of one neutron of a nn-pair. The deviation $\delta = 0.028$ fm from fitting the data of the isotopes with $Z \geq 24$ of Table IIIA [5]. For the experimental radii of the 147 isotopes with $Z \geq 6$ of the table $\delta = 0.037$ fm.

One should notice here that the radius of a core α - cluster $4^{1/3} r_{p/n}$ is not its mass radius. It rather defines the space which can't be occupied by excess neutrons, because there is some structure made of pn-pairs. The results of fitting experimental radii show that the charge radius of an α-cluster slowly changes with growing Z (otherwise δ would not depend of the group of data selected for fitting). So does the α- cluster's body radius.

The real mass radii of all nucleons considered as elementary bricks of a nucleus should be equal and they can not change, because the binding energies are too small in comparison with their masses. The well known formula $R = r_0 A^{1/3}$ of the liquid drop model with the fitted value r_0 =0.95 fm for the isotopes with $Z \geq 24$ of the Table IIIA [5] gives the considerably bigger deviation $\delta = 0.067$ fm and for 147 isotopes with $Z \geq 6$ rms deviation is $\delta = 0.12$ fm, which is considerably worse than those of the equations (9-12). This clearly shows that the distances between nucleons in a nucleus are not related to their mass radii.

The experimental data of the other isotopes with $60 \leq Z, Z_1 \leq 77$ are presented in [5] in a separate table, Table IIIC, because they are treated as deformed ones. The radii for the isotopes calculated by (9-12) with the N_α^{ml} obtained in the analysis of experimental binding energies (see section 2) are in an agreement with the data in the table, although they are always less than the experimental radii on several hundredths of fm.

2 Binding energies and radii of the isotopes with $N \geq Z$

For the nuclei with $Z, Z_1 \geq 30$ the number N_α^{ml} is found from the analysis of experimental binding energies [6] with average deviation of calculated energies from experimental values in 2.5 MeV. Some examples are given in the table in Appendix. All isotopes with growing the number ΔN have the N_α^{core} increasing and N_α^{ml} correspondingly decreasing. At β-stable isotopes N_α^{ml} reaches the value 2 for some of the isotopes with $Z = 38 \div 43$ and $N^{ml} = 4$ for some of the isotopes with $66 \leq Z, Z_1 \leq 79$. For the other β-stable isotopes $N_\alpha^{ml}=$ 3. The β-stable isotopes with $Z, Z_1 \geq 85$ have $N_\alpha^{ml} = 4, 5$. The calculated binding energies of the isotopes with $12 \leq Z, Z_1 \leq 29$ do not depend much on N_α^{ml} varied within $2 \div 5$. So for the β-stable isotopes of the nuclei with $24 \leq Z, Z_1 \leq 29$ it is suggested that N_α^{ml} equals 3 as it does for the heavier nuclei. For the light nuclei having core $12 \leq Z, Z_1 \leq 23$ the calculations on (10-12) with the value $N_\alpha^{ml} = 5$ give a small deviation δ from the experimental radii. Spreading the common rule that with growing A the value N_α^{ml} decreases to 3 for these nuclei makes the deviations smaller.

The specific density of core binding energy is calculated as $\rho = E_{core}/(N_\alpha^{core} v_\alpha)$ where E_{core} is (4) for the isotopes with $Z, Z_1 \geq 30$, $v_\alpha = 4/3\pi r_\alpha^3$ at the charge radius of the core α - cluster $r_\alpha = 1.595$ fm taken from fitting the experimental radii by (9) and (11). The value ρ grows slowly with the number of excess neutrons and at β-stable isotopes it reaches a saturation value $\rho \approx 2.6 \pm 0.1$ MeV/fm^3. For example, the β-stable isotopes with $Z = 30$, 60 and 80 have N^{ml}= 3 and $\rho = 2.48 \div 2.67$ MeV/fm^3, $2.53 \div 2.63$ MeV/fm^3 and $2.47 \div 2.55$ MeV/fm^3 correspondingly. Then the narrow strip of binding energies of β - stability is to be outlined by E (2) with $E_{core} = N_\alpha^{core} v_\alpha \rho$. With N_α^{ml} =3 (for $Z, Z_1 \leq 84$) and $\rho = 2.55$ MeV/fm^3 the binding energy E (2) is a function of one variable N_α (Z)

$$E = E_{N_\alpha^{ml}} + N_\alpha^{core} v_\alpha \rho - 2N_\alpha^{ml} 2N_\alpha^{core} e^2/R_\alpha, \tag{13}$$

where $N_\alpha^{core} = N_\alpha - N_\alpha^{ml}$ and $R_\alpha = 2.168(N_\alpha - 4)^{1/3}$ (for the $Z \leq 22$ the empirical values R_α are used, see (17) in [3]). In Fig. 1 the graph of the function (13) is given in comparison with the experimental values of the lightest and the



Fig. 1. The binding energy of beta-stable isotopes. Dots indicate the experimental energies of the lightest and the heaviest even beta-stable isotopes. Two lines indicate the function (13) with N_α^{ml}= 3 and 5.

heaviest even Z beta-stable isotopes depicting the boundaries of β-stability. The width of the strip is determined by the variation of ρ within $2.5 \div 2.7$ MeV/fm^3 and by the number N_α^{core}. The physical meaning of $v_\alpha \rho$=43.34 MeV is the core binding energy per one α - cluster. Then the number of excess neutrons of β - stability is defined by (5) and by the eq. $E_{\Delta N} = 43.34 N_\alpha^{core} - E_{N_\alpha^{core}} - 6\epsilon_{\alpha\alpha}$.

As it was shown in [1] the binding energies of the excess nn-pairs in β-stable nuclei obey the eq. (5) with the accuracy in a few MeV. Therefore the energy E (2,8) with $E_{\Delta N}$ (5) can be calculated only for those isotopes that have the excess neutrons inside the core, i.e. for the $A \leq A^{st}$ where A^{st} is the mass number of the heaviest β-stable isotope among those belonging to one Z. Examples are given in Table 1 (see Appendix) for the isotopes with Z = 30, 31, 80, 81. For $Z = 30$ and 31 all the isotopes with $N \geq Z$ with known energies are given. One can see that for the isotopes with $A > A^{st}$ the values of the nn-pair separation energy $E_{N_{nn}}^{sep} = E_{A'(Z,N+2)} - E_{A(Z,N)}$ less than $E_{N_{nn}}$ (5). It allows one to suggest that the nn-pairs come out of the core not affecting the core density. In such cases the binding energy is equal to the sum of E (2) for the isotope A^{st} and $\sum E_{i_{nn}}^{sep}$ of the last nn-pairs.

One can see that $E_{N_{nn}}^{sep}$ for both isotopes $A(Z,N)$ and $A_1(Z_1, N+2)$ in most of the cases are almost equal. It approves the idea that the cores of the nuclei with Z and Z_1 are mostly equal. One can see from Table 1, that the separation energies $E_{N_{nn}}^{sep}$ for the most of the pairs of $A(Z,N)$ and $A_1(Z_1, N+2)$ differ within 1 MeV. It allows one to make a prediction for some isotopes with $A > A^{st}$, when the energy of the complementary nucleus has been already measured. Some of the predicted values are $E_{123(45,78)}$ =1007 MeV, $E_{126(46,80)}$ = 1034 MeV, $E_{139(51,88)}$= 1132 MeV, $E_{145(53,92)}$ = 1171 MeV, $E_{148(54,94)}$ = 1193 MeV, $E_{189(73,116)}$=1500 MeV, $E_{193(75,118)}$=1529 MeV, $E_{175(81,94)}$=1348 MeV, $E_{261(101,160)}$=1929 MeV.

In accordance with the representation of a core as a liquid drop one can suggest that the excess nn-pairs concentrate inside the core near its surface due to the surface tension, so that the filling out the space inside the core goes from surface to center. The very first nn-pairs can be at the boundary of the core defined by core's charge radius. Therefore for the isotopes lighter than stable ones the equations (9-11) should predict the charge radii. For the β - stable isotopes and the heavier ones with the nn-pairs out of the core one should use (12) to estimate the nuclear size. Then the nuclear radii with growing A should first decrease with decreasing N_α^{ml} till the stable isotopes, which have approximately equal radii calculated by (9-12). For the heavier isotopes the radii slowly increase (12) with growing A because of the growing amount of nn-pairs on the surface of the core.

References

[1] G. K. Nie, arXiv:0707.4291v3 [nucl-th] 20Sep2007

[2] G. K. Nie, *Mod. Phys. Lett. A*, **21**, 1889 (2006).

[3] G. K. Nie, *Mod. Phys. Lett. A*, **22**, 227 (2007).

[4] P. D. Norman, *Eur. J. Phys.* **14**, 36 (1993).

[5] G. Fricke et al, *Atomic Data and Nuclear Data Tables* **60**, 207 (1995)

[6] CDFE online service, http://cdfe.sinp.msu.ru/

3 Appendix

Table 1. Binding energies and radii of the isotopes with Z=30, 31, 80, 81. ΔN is the number of excess neutrons in the core. E_{exp} is from [6], E_{nn}^{sep} is the separation energy of the last nn-pair, N_α^{ml} is the number of alpha-clusters out of the core, R_{exp} is from [5]. * indicates a β-stable nucleus

Z	ΔN	A	E_{exp}	E_{nn}^{sep}	N_α^{ml}	E (2,8)	R_{exp}	$R(12)$	$R(11)$	$R(10)$	$R(9)$
			MeV	MeV		MeV	fm	fm	fm	fm	fm
30	0	60	515	0	3	514		3.841	3.914	3.954	3.934
31	0	63	541	0	3	538		3.897	3.949	4.006	3.977
30	2	62	538	23	3	535		3.864	3.914	3.954	3.934
31	2	65	563	22	3	559		3.919	3.949	4.006	3.977

Z	ΔN	A	E_{exp}	E_{nn}^{sep}	N_{α}^{ml}	E (2,8)	R_{exp}	$R(12)$	$R(11)$	$R(10)$	$R(9)$
			MeV	MeV		MeV	fm	fm	fm	fm	fm
30	4	*64	559	21	3	556	3.928	3.886	3.914	3.954	3.934
31	4	67	583	20	3	579		3.941	3.949	4.006	3.977
30	6	*66	578	19	3	575	3.948	3.908	3.914	3.954	3.934
31	6	*69	602	19	3	599	3.996	3.962	3.949	4.006	3.977
30	8	*68	595	17	3	594	3.965	3.930	3.914	3.954	3.934
31	8	*71	619	17	3	618	4.011	3.984	3.949	4.006	3.977
30	10	70	610	16	3	613	3.983	3.952	3.914	3.954	3.934
31	10	73	635	16	3	636		4.005	3.949	4.006	3.977
30	12	72	626	16	3			3.973	3.914	3.954	3.934
31	12	75	650	16	3			4.025	3.949	4.006	3.977
30	14	74	640	14	3			3.995	3.914	3.954	3.934
31	14	77	663	13	3			4.046	3.949	4.006	3.977
30	16	76	652	13	3			4.015	3.914	3.954	3.934
31	16	79	676	13	3			4.067	3.949	4.006	3.977
30	18	78	663	11	3			4.036	3.914	3.954	3.934
31	18	81	688	12	3			4.087	3.949	4.006	3.977
30	20	80	674	11	3			4.057	3.914	3.954	3.934
31	20	83	695	7	3			4.107	3.949	4.006	3.977
30	22	82	681	7	3			4.077	3.914	3.954	3.934
31	22	85	702	7	3			4.127	3.949	4.006	3.977
80	12	172	1327		3+3	1327		5.362	5.512	5.458	5.455
81	12	175			3+3	1349		5.391	5.536	5.486	5.477
80	14	174	1348	21	3+3	1343		5.373	5.512	5.458	5.455
81	14	177	1369		3+3	1367		5.385	5.522	5.474	5.477
80	16	176	1370	22	2+3	1369		5.367	5.498	5.446	5.455
81	16	179	1390	21	2+3	1392		5.396	5.522	5.474	5.477
80	18	178	1390	20	2+3	1386		5.379	5.498	5.446	5.455
81	18	181	1410	20	2+3	1409		5.408	5.522	5.474	5.477

Z	ΔN	A	E_{exp} MeV	E_{nn}^{sep} MeV	N_{α}^{ml}	$E(2{,}8)$ MeV	R_{exp} fm	$R(12)$ fm	$R(11)$ fm	$R(10)$ fm	$R(9)$ fm
80	20	180	1410	20	1+4	1411		5.391	5.498	5.446	5.455
81	20	183	1430	20	1+4	1431		5.419	5.522	5.474	5.477
80	22	182	1430	20	1+4	1427		5.402	5.498	5.446	5.455
81	22	185	1450	20	1+4	1448		5.431	5.522	5.474	5.477
80	24	184	1449	20	1+3	1450		5.396	5.485	5.433	5.455
81	24	187	1468	18	1+4	1463		5.425	5.509	5.461	5.477
80	26	186	1467	18	1+3	1466		5.408	5.485	5.433	5.455
81	26	189	1487	19	1+3	1489		5.436	5.509	5.461	5.477
80	28	188	1485	18	1+3	1480		5.419	5.485	5.433	5.455
81	28	191	1504	17	1+3	1504		5.447	5.509	5.461	5.477
80	30	190	1502	17	4	1505		5.431	5.485	5.433	5.455
81	30	193	1526	22	4	1525		5.459	5.509	5.461	5.477
80	32	192	1519	17	4	1519		5.442	5.485	5.433	5.455
81	32	195	1539	13	5	1536		5.487	5.522	5.474	5.477
80	34	194	1535	16	4	1533		5.453	5.485	5.433	5.455
81	34	197	1555	16	4	1553		5.481	5.509	5.461	5.477
80	36	*196	1551	16	3	1554		5.448	5.475	5.421	5.455
81	36	199	1571	16	4	1566		5.492	5.509	5.461	5.477
80	38	*198	1566	15	3	1567	5.448	5.459	5.475	5.421	5.455
81	38	201	1586	15	3	1590		5.487	5.499	5.449	5.477
80	40	*200	1581	15	3	1580	5.457	5.470	5.475	5.421	5.455
81	40	*203	1601	15	3	1603	5.472	5.498	5.499	5.449	5.477
80	42	*202	1595	14	3	1592	5.467	5.481	5.475	5.421	5.455
81	42	*205	1615	14	3	1615	5.483	5.509	5.499	5.449	5.477
80	44	*204	1609	14	3	1604	5.478	5.492	5.475	5.421	5.455
81	44	207	1628	13	3	1627		5.520	5.499	5.449	5.477